

Comment on “Energy Dependence of Electron Lifetime in Graphite Observed with Femtosecond Photoemission Spectroscopy”

In a recent letter, Xu *et al.* [1] reported measurement of electron lifetime in graphite with photoemission spectroscopy. By performing a least-square fitting of the observed energy dependence of the electron lifetime and adapting a layered two-dimensional model for graphite, the authors attributed the measured electron-electron scattering rate to contributions due to acoustic plasmon excitations. They came to the conclusion that for a layered two-dimensional electron gas with layer separation small compared with the average inter-electron distance within the layer, the acoustic plasmon excitations dominate electron-electron scattering processes and give rise to a linear energy dependence of the scattering rate for electrons with energies close to the Fermi energy. The authors further implied that the scattering due to acoustic plasmon excitations may also be applied to explain the linear energy and temperature dependence of carrier lifetimes observed in the normal state of high- T_c superconductors since most of the high- T_c superconductor materials have similar layered electronic structures. The purpose of this comment is to point out that these authors’ theoretical interpretations of their experimental data and, most importantly, the suggested implication for high- T_c superconductors are incorrect and misleading.

Electron-electron scattering contains contributions from both single particle electron-hole excitations and collective plasmon excitations. In a layered electronic system, the plasmon spectrum consists of optical and acoustic modes [2]. The scattering rate due to the acoustic plasmon excitations can be expressed analytically in a narrow energy window as [1,3]

$$\frac{1}{\tau_{ac}(k)} = C(\sqrt{\epsilon_k} - \sqrt{\epsilon_c})^2, \quad (1)$$

where C is a constant, $\epsilon_k = \hbar^2 k^2 / 2m$ is electron energy, and ϵ_c is a threshold energy. The above expression is valid only for electrons with energies $\epsilon_c < \epsilon_k < \epsilon^*$ [3]: for electrons with energies smaller than ϵ_c , plasmon excitations do not contribute to electron scattering because of restrictions from momentum and energy conservations; for electrons with energies higher than ϵ^* , $\tau_{ac}(k)^{-1}$ actually decreases as a function of ϵ_k , and Eq.(1) does not apply.

Since the threshold energy ϵ_c is higher than the Fermi energy and decreases with decreasing layer separation [3], the acoustic plasmons do not contribute to scattering of electrons with energies close to the Fermi surface, except for extremely small layer separations where ϵ_c is sufficiently close to ϵ_F . In this small layer separation limit, Eq.(1) gives $\tau_{ac}(k)^{-1} \sim (\epsilon_k - \epsilon_F)^2$ for $\epsilon_k \rightarrow \epsilon_c \simeq \epsilon_F$, which is quadratic, not linear, in energy. We conclude, therefore, that in sharp contrast to statements made in ref. [1], the electronic scattering rate due to acoustic plasmon excitations does not show linear energy dependence under any circumstances. The theoretical conclusions of ref. [1] were obtained by comparing the measured data with Eq.(1), which, as should be obvious from our comment, does not show a linear $(\epsilon - \epsilon_F)$ behavior for any values of electron energy. An additional problem is that the experimental data, with electron energies on the order of eV , are in the high electron energy regime where Eq.(1) is not valid. It is then clear that the theoretical interpretations of ref. [1] are wrong.

There is another problem associated with attributing the measured scattering rate solely to contributions due to acoustic plasmon excitations. In a layered electronic gas, the spectral weight of acoustic plasmons decreases as the layer separation decreases, since only the optical plasmon mode survives in the zero layer separation limit. In graphite, the layer separation is $3.7\text{\AA}$ and the average inter-electron distance within the layer is on the order of $100 \sim 200\text{\AA}$. Under this small layer separation condition, contributions to the scattering rate from excitations other than the acoustic plasmons may not necessarily be negligible. The authors of ref. [1] uncritically disregarded other possible sources of contributions without any quantitative examination.

We emphasize that the implication of ref. [1] for carrier lifetimes in the normal state of high- T_c superconductors, as suggested by the authors, is invalid. Many experiments on the normal state of high- T_c superconductor materials suggest linear energy and temperature dependence of lifetimes for carriers close to the Fermi energy. It is well known that the anomalous linear energy and temperature dependence of the lifetime is not expected from a Fermi-liquid type many-body theory for a single-layer uniform two-dimensional electron gas. In fact, the standard Fermi-liquid result for the scattering rate of a two-dimensional

electron gas is an $(\epsilon - \epsilon_F)^2 \ln(\epsilon - \epsilon_F)$ behavior for ϵ close to ϵ_F , which is extremely well-verified experimentally [4]. From the discussion above, it is clear that acoustic plasmon contributions in a layered two-dimensional electron gas do not produce a linear $(\epsilon - \epsilon_F)$ behavior for the scattering rate either. It is well-established [5] that both two- and three-dimensional electron systems, at least within the standard perturbative many-body picture, have scattering rates which are essentially quadratic in energy, *i.e.* $(\epsilon - \epsilon_F)^2$ within logarithmic corrections, for electrons close to the Fermi surface. The same is also true for a layered electron gas system, at least within the standard Fermi liquid many-body perturbation theory.

Lian Zheng and S. Das Sarma

Department of Physics

University of Maryland

College Park, Maryland 20742-2111

PACS numbers: 73.20Dx, 73.20.Mf, 73.61.-r

REFERENCES

- [1] S. Xu *et al.*, Phys. Rev. Lett. **76**, 483 (1996).
- [2] S. Das Sarma and A. Madhukar, Phys. Rev. B. **23**, 805 (1981); S. Das Sarma and J.J. Quinn, *ibid.* **25**, 7603 (1982).
- [3] P. Hawrylak, G. Eliasson, and J.J. Quinn, Phys. Rev. B. **37**, 10187 (1988).
- [4] L. Zheng and S. Das Sarma, Phys. Rev. B. **53**, (15 April, 1996); S.Q. Murphy *et al.*, Phys. Rev. B. **52**, 14825 (1995).
- [5] R. Shankar, Rev. Mod. Phys. **66**, 129 (1994) and references therein.